

ARRHYTHMIAS AFTER CARDIAC SURGERY-A PROSPECTIVE STUDY IN THE INSTITUTE OF CARDIOVASCULAR DISEASES OF TÂRGU-MUREȘ

MATHE ZSOMBOR¹, HORAȚIU SUCIU², RADULY GERGO³

^{1,3} PhD candidate University of Medicine and Pharmacy Târgu-Mureș, ²University of Medicine and Pharmacy Târgu-Mureș

Keywords: surgery, cardiac, arrhythmias

Abstract: Introduction: Cardiac arrhythmias are the most common complication in cardiac surgery and they play a major role in postoperative mortality and morbidity. New-onset of atrial fibrillation is the most common arrhythmic complication after coronary artery bypass graft surgery and it may result in haemodynamic compromise, thrombotic events, anxiety, and thus a prolonged hospitalization. Materials and methods: A prospective study was performed in the Institute of Cardiovascular Diseases and Transplantation of Targu Mures, Romania. We collected data between January and June 2017. For statistical analysis T-test and Fisher-test (GraphPad Prism 6.0 software, GraphPad Software, Inc, La Jolla, CA 92037 USA) were used. Significance level was set at $p < 0.005$. Results: In the arrhythmic group, the mean age was higher than in the control group. The ejection fraction was lower in the arrhythmic group. In the arrhythmic group, more cases needed higher doses of inotropic drugs than in the control group. The arrhythmic group presented statistically significant hemodynamic and renal postoperative complications. Discussions: Cardiac dilatation, myocardial atrophy, decrease of conduction tissue and atrial fibrosis caused by aging may be responsible for the increased risk of atrial fibrillation after CABG. Echography findings that suggested the increased risk of atrial fibrillation in the postoperative period was the enlarged atrium. The blood supply to the right atrium, the sinoatrial node and the atrioventricular node is mainly conveyed via the right coronary artery, thus the right coronary artery stenosis may be a predictor. The statistically significant hemodynamic and renal postoperative complications in our study are among the most common postoperative complications described in the literature. Conclusions: Postoperative atrial fibrillation is more frequent in the elderly patients (above 60 years) with increased left atrium. Other disturbances that predispose to this arrhythmia are hypoxia and metabolic acidosis.

INTRODUCTION

Post-operative arrhythmias are frequent complications in patients undergoing cardiac surgery. Causing a significant increase in post-operative morbidity and mortality, arrhythmias are likely to be present in normal patients.(1) Still, they can cause hemodynamic instability in the recovery period after a cardiac surgery.(2,3)

New-onset postoperative atrial fibrillation (POAF) is a common arrhythmic complication after first-time surgical myocardial revascularization.(4,5) Various studies reported an incidence between 11-40%.(6,7)

POAF can cause haemodynamic compromise, acute kidney injuries, long term thromboembolic events along prolonged hospitalization, increased in-hospital and long-time mortality in comparative studies presenting results from patients in sinus rhythm after cardiac surgery.(8,9,10,11)

It may be related to clinical and perioperative factors.(12)

Among a multitude of studies, in 2004, Mathew and colab. confirmed that age over 70 years is associated with an increased risk of atrial fibrillation (AF).(7,13,14)

Left atrial enlargement, diastolic dysfunction and left ventricular hypertrophy are considered factors in POAF occurrence.(15)

Cardiopulmonary by-pass time (CBP), atrial and venous cannulation, myocardial ischemia, volume and

electrolyte imbalances, hypotension, are also considered to represent a cause for the increased number of POAF.(13,15)

Likewise, the hyperadrenergic condition following surgical stress and inotropic drugs contribute to postoperative atrial fibrillation.(16)

AF may be prevented by focussing mainly on anti-arrhythmic medication: digitalis, beta-blockers, calcium channel blockers and amiodarone. The aims of treatment for post-operative AF are control of the ventricular response, prevention of cardioembolic events and restoring normal sinus rhythm.(17)

MATERIALS AND METHODS

The present prospective study used the medical records database from the Cardiac Surgery Clinic, Institute of Cardiovascular Diseases and Transplantation, Târgu-Mureș, Romania. Secondary to obtaining the ethical committee approval, we studied all patients' files that underwent heart surgery between January and June 2017. All investigations were conducted according to the standards of the declaration of Helsinki.

Patients' data were included based on the following criteria: unrestricted access to patients' medical records, existence of clinical, imagistic data, laboratory results and perioperative Intensive Care Unit (ICU) data.

Exclusion criteria consisted of previous cardiac surgery, chronic kidney disease (CKD), thyroid disorders,

³Corresponding author: Raduly Gergo, Str. Ghe. Marinescu, Nr. 38, Târgu-Mureș, România, E-mail: raduly_gergo@yahoo.com, Phone: +0747 714387

Article received on 22.07.2018 and accepted for publication on 03.09.2018
ACTA MEDICA TRANSILVANICA September 2018;23(3):31-34

CLINICAL ASPECTS

history of supraventricular arrhythmia. A number of 90 consecutive patients undergoing first-time cardiac surgery were included.

Our statistical analysis did not include the number of deaths during the procedures nor the 24-hours-after-surgery deaths.

We followed the presence of newly emerged AF based on gender, age, surgery type (isolated CABG +/- valve replacement), preoperative comorbidities, surgery time, echocardiographic parameters (ejection fraction-EF and left atrium size-LAS), post-surgical organ dysfunctions (neurologic, respiratory, renal, gastro-intestinal injuries) and laboratory results (electrolytes, complete blood count, creatinine and Astrup parameters – pH, SaO₂), cardio-pulmonary by-pass and total ischemic times.

Ejection fraction (modified Simpson's rule) and left atrial size were assessed by 2D echocardiography. Classic cut-off values were set at 45% for EF and 45mm for LAS.(18)

For the statistical analysis we used the T-test and Fisher's-test (statistical significance p<0.05), using GraphPad Prism 6.0 software, GraphPad Software, Inc, La Jolla, CA 92037 USA.

RESULTS

All patients (66 males, 24 females) were divided into two groups. Patients presenting arrhythmia (38/90 patients - 42.20%) - the arrhythmic group was compared with a control group (52/90pts - 57.70%) - patients without AF.

Among all participants (73.33% male patients and 26.66% female patients) the mean age was 60.33 years, still there was no statistical significance (p>0.05); in the arrhythmic group, mean age was higher compared to the control group.

We also studied the preoperative comorbidities, such as hemodynamic disturbances, respiratory-, renal-, gastrointestinal disorders and presence of diabetes mellitus. No statistically significant differences between the two groups were observed (table no. 1).

Table no. 1. Preoperative comorbidities

Variables	Arrhythmic group (n=38)	Control group (n=52)	p value	OR, CI
Hemodynamic disturbances				
Respiratory disorders	12	18	p=0.823	OR:0.87, 95% CI:0.36 to 2.17
CKD	10	10	p=0.451	OR:1.5, 95% CI: 0.589 to 3.815
Diabetes mellitus	6	8	p=0.999	OR 1.03, 95%CI:0.307 to 3.052
Other	32	38	p=0.305	OR 1.96, 95%CI:0.6694 to 5.817

n-number of cases; CKD-chronic kidney disease; OR- odds ratio, CI- confidence interval.


The arrhythmic group presented a reduced ejection fraction, not statistically significant (p>0.05) compared to the control group. When analysing the echocardiographic data, a statistically significant (p<0.05) difference between the left atrium size was estimated within the two groups (table no. 2).

Table no. 2. Echocardiographic findings

Variables	Arrhythmic group	Control group	p value
LVEF (mean)	51.53%	53.67%	0.3935
LA size (mean)	47.50 mm	43.05 mm	0.0538


LVEF-left ventricular ejection fraction; LA- left atrium.

Figure no. 1. Type of surgery



As the type of surgery, 26 patients from the control group underwent coronary artery bypass surgery (CABG), 22 of the control group underwent valve replacement surgery and 4 underwent both (figure no. 1).

Figure no. 2. Cardio-pulmonary by-pass and miocardic ischaemia times



In the arrhythmic group the cardio-pulmonary bypass time was higher than in the control group, but the miocardic ischaemia time was higher in the control group (figure no. 2).

Within the two groups Na⁺, K⁺ and SaO₂ daily levels were computed. For arterial blood oxygen saturation, a statistically significant (p=0.0001) difference was estimated. When determining the difference between the electrolytes levels, for K⁺ there was a statistically significant (p=0.0014) disparity. The creatinine and white blood cell levels within the arrhythmic group were higher, with a statistically significant (p=0.005) value for the creatinine level. For the haematocrit, haemoglobin and pH, lower levels were documented among the arrhythmic group patients, yet not statistically significant (p>0.05) (table no. 3.).

Table no. 3. Paraclinical parameters - mean values

Variables mean values	Arrhythmic group	Control group	p value
Na ⁺ (mEq/L)	141.8	141.6	0.6397
K ⁺ (mEq/L)	4.021	3.819	0.0014
SaO ₂ %	97.72	98.89	0.0001
Creatinine (mg/dl)	1.54	1.081	0.0050
WBC (/mm ³)	11790	9986	0.0633
Hematocrit (%)	29.68	30.22	0.4247
Hemoglobin (mmol/L)	9.4	9.7	0.1664
pH	7.37	7.39	0.0732

Na⁺ - sodium plasma level, K⁺ - Potassium plasma level, WBC - white blood cells, SaO₂ - arterial oxygen saturation.

The arrhythmic group presented statistically significant (p=0.0063) hemodynamic and renal postoperative complications (table no. 4.).

A higher number of patients included in the arrhythmic group required high doses of inotropic drugs, in comparison to the control group, statistically significant for

CLINICAL ASPECTS

Dobutamine ($p=0.0204$) and Noradrenaline ($p=0.0289$) (table no. 5).

Table no. 4. Postoperative complications

Postoperative complication	Arrhythmic group (n=38)	Control group (n=52)	p value
Neurologic	0	0	1
Respiratory	2	6	0.4594
Renal	6	0	0.0044
Gastrointestinal	6	2	0.0660
Hemodynamic	20/18	6/46	0.0063
Death	2	0	0.1755

n=number of cases;

Table no. 5. Drugs

	Arrhythmic group (n=38)	Control group (n=52)	p value
Dopamine (> 5 µg/kg/min)	12	8	$p=0.0680$
Dobutamine (> 5 µg/kg/min)	10	4	$p=0.0204$
Adrenaline	4	2	$p=0.2363$
Noradrenaline	4	0	$p=0.0289$

n=number of cases

and diabetes mellitus, the literature mentions some comorbidities as predictive factors for atrial fibrillation. In hypertension there is increased fibrosis and dispersion of atrial refractoriness. The blood supply to the right atrium, the sinoatrial node and the atrioventricular node is mainly from the right coronary artery, thus stenosis of the right coronary artery may be a predictor. Chronic obstructive lung disease patients frequently have premature atrial contractions, which may trigger the initiation of atrial fibrillation.(20,22)

In terms of laboratory parameters, we found higher Na, K levels, but these were not significant statistically. We found increased leukocyte levels in our patients with postoperative atrial fibrillation. The inflammatory processes initiated by extracorporeal circulation (ECC) might alter intra atrial conduction and other electrophysiological properties of the atria, leading to postoperative AF.(27)

The statistically significant hemodynamic and renal postoperative complications in our study are among the most common postoperative complications described in the literature: myocardial infarction, persistent congestive heart failure, respiratory failure, various infectious complications, renal failure, severe hypotension and shock, multisystemic failure, and cardiopulmonary arrest.(19) Postoperative pneumonia and mechanical ventilation longer than 24 hours have been shown to be independent postoperative predictors of AF, as well as atrial pacing, and need of postoperative inotropic agents.(20)

DISCUSSIONS

Atrial fibrillation is one of the most common postoperative complications in cardiac surgery, prevalence depending on the type of surgery, population profile, detection methods. Its incidence after elective coronary-aortic bypass graft surgery varies in a wide range, from region to region.(19)

Postoperative atrial fibrillation may be asymptomatic, but symptoms include palpitations, breathlessness, chest pain, sweating, and hypotension.(19)

The postoperative arrhythmia episodes are diagnosed by continuous monitoring and the use of 12 lead electrocardiogram. Episodes occur within the first 6 days following cardiac surgery.(19)

In our study, we found atrial fibrillation mostly in elderly men, as age is an independent risk factor.(20,21,22) Cardiac dilatation, myocardial atrophy, decrease of conduction tissue and atrial fibrosis caused by aging may be responsible for the increased risk of atrial fibrillation after CABG. The difference between genders may be explained by the sex differences in ion-channel expression and hormonal effects on autonomic tone.(20)

Echography findings that suggested the increased risk of atrial fibrillation in the postoperative period was the enlarged atrium. Many studies described the enlarged atrium as potent and independent predictor of atrial fibrillation in the postoperative period.(19,21)

There is a correlation between the prolongation of the by-pass time and the presence of postoperative arrhythmias. In this prolonged time, the myocyte membrane becomes sensitive, the action potential and the refractory period decrease, which predisposes to activation of pacemaker sites and re-entry cycles.(23,24)

Atrial fibrillation appears more frequently after valve surgeries, which is explained by the increase of atrium volume and presence of the surgical cicatrix.(24,25,26)

In our study, the ejection fraction was lower in the arrhythmic group, but not statistically significant, however, it is mentioned in the literature as a predicting factor.(22)

Although we did not find statistically significant changes among comorbidities such as hemodynamic disturbances, respiratory-, renal-, gastrointestinal disturbances

CONCLUSIONS

We can conclude that based on our study, postoperative atrial fibrillation is more frequent in elderly patients (above 60 years) with increased left atrium. Other disturbances that predispose to this arrhythmia are hypoxia and metabolic acidosis. However, we must not forget the increased number of leucocytes due to the systemic inflammatory response, anemia and the need of high level of inotropic drugs, but in these cases the changes were not statistically significant.

REFERENCES

- Ahlsson A, Bodin L, Fengsrud E, et al: Patients with postoperative atrial fibrillation have a doubled cardiovascular mortality. *Scand Cardiovasc. J* 2009;43(5):330-6.
- Albert M Ai, Halevy Ni, Antman E Mi. Preoperative evaluation for Cardiac Surgery Cohn Lh, ed. *Cardiac Surgery in the Adult*. New York: McGraw-Hill. 2008; p. 261-280.
- Crystal E, Connolly SJ, Sleik K, Ginger TJ, Yusuf S. Interventions on prevention of postoperative atrial fibrillation in patients undergoing heart surgery: a meta-analysis. *Circulation*. 2002;106(1):75-80.
- Turk T, Vural H, Eris C, Ata Y, Yavuz S. Atrial fibrillation after off-pump coronary artery surgery: a prospective, matched study. *J Int Med Res*. 2007;35(1):134-42.
- Hakala T, Pitkänen O, Hippeläinen M. Feasibility of predicting the risk of atrial fibrillation after coronary artery bypass surgery with logistic regression model. *Scand J Surg*. 2002;91(4):339-44.
- Valkey AJ, Benjamin EJ, Lubitz SA. New-onset atrial fibrillation during hospitalisation. *J Am Col Cardiol*. 2014;64(22):2432-3.
- Mathew JP, Fontes ML, Tudor IC, et al. Investigators of the ISCHEMIA research and education foundation; multicentre study of perioperative ischemia research group. A multicentre risk index for atrial fibrillation after cardiac surgery. *JAMA*. 2004;291(14):1720-9.

CLINICAL ASPECTS

8. Saxena A, Dinh DT, Smith JA, Shardey GC, Reid CM, Newcomb AE. Usefulness of postoperative atrial fibrillation as an independent predictor for worse early and late outcomes after isolated coronary artery bypass grafting (multicenter Australian study of 19,497 patients). *Am J Cardiol.* 2012;109(2):219-225.
9. Helgadóttir S, Sigurdsson MI, Ingvarsdóttir IL, Arnar DO, Gudbjartsson T. Atrial fibrillation following cardiac surgery: risk analysis and long-term survival. *J Cardiothorac Surg.* 2012;7(87):1-7.
10. Phan K, Ha HS, Phan S, Medi C, Thomas SP, Yan TD. New-onset atrial fibrillation following coronary bypass surgery predicts long-term mortality: a systematic review and meta-analysis. *Eur J Cardiothorac Surg.* 2015;48(6):817-824.
11. Hravnak M, Hoffman LA, Saul MI, et al: Resourceutilizationrelatedtoatrialfibrillationaftercoronaryartery bypass grafting. *Am J Crit Care.* 2002;11(3):228-8.
12. El-Chami MF, Kilgo P, Thourani V, et al. New-onset atrial fibrillation predicts long-term mortality after coronary artery bypass graft. *J Am Coll Cardiol.* 2010;55(13):1370-1376.
13. Aranki SF, Shaw DP, Adams DH, et al. Predictors of atrial fibrillation after coronary artery surgery. Current trends and impact on hospital resources. *Circulation.* 1996;94:390-7.
14. Mathew JP, Parks R, Savino JS, et al. Atrial fibrillation following coronary artery bypass graft surgery: predictors, outcomes, and resource utilization. MultiCenterStudy of Perioperative Ischemia Re- search Group. *JAMA* 1996;276:300-6.
15. Echahidi N, Pibarot P, O'Hara G, Mathieu P. Mechanisms, prevention, and treatment of atrial fibrillation after cardiac surgery. *J Am CollCardiol.* 2008;51(8):793-801.
16. Kalman JM, Munawar M, Howes LG, et al. Atrial fibrillation after coronary artery bypass grafting as associated with sympathetic activation. *Ann ThoracSurg.* 1995;60(6):1709-1715.
17. Kirchhof P, Benussi S, Kotecha D, Ahlsson A, Atar D, Casadei B. 2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Top of Form Eur Heart J.* 2016;37(38):2893-2962. Bottom of Form
18. Raitt MH, Volgman AS, Zoble RG, CharbonneauLet, Pader FA, O'Hara GE, et al. Prediction of the recurrence of atrial fibrillation after cardioversion in the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study. *Am Heart J.* 2006;151(2):390-6.
19. Kaireviciute D, Aidietis A, Lip GY .Atrial fibrillation following cardiac surgery: clinical features and preventative strategies. *Eur Heart J.* 2009;30(4):410-25.
20. Jung W, Meyerfeldt U, Birkemeyer R - Atrial arrhythmias after cardiac surgery in patients with diabetes mellitus. *Clin Res Cardiol.* 2006 Jan;95 Suppl1:i88-97.
21. Hayashida N, Shojima T, Yokokura Y, Hori H, Yoshikawa K, Tomoeda H, Aoyagi S. - P-wave signal-averaged electrocardiogram for predicting atrial arrhythmia after cardiac surgery. *Ann Thorac Surg.* 2005 Mar;79(3):859-64.
22. Karaca M, Demirbas MI, Biceroglu S, Cevik A, Cetin Y, Arpaz M, Yilmaz H - Prediction of early postoperative atrial fibrillation after cardiac surgery: is it possible? *Cardiovasc J Afr.* 2012 Feb;23(1):34-6.
23. Nair SG, et al. Atrial fibrillation after cardiac surgery. *Ann Card Anaesth* 2010, 13: 196-205.
24. Patel D, Gillinov MA, Natale A, et al. Atrial Fibrillation after Cardiac Surgery: Where are we now? *Indian Pacing and Electrophysiology Journal.* 2008;8:281-291.
25. Echahidi N, Pibarot P, O'Hara G, et al. Mechanism, Prevention, and Treatment of Atrial Fibrillation after Cardiac Surgery. *Journal of the American College of Cardiology.* 2008;51:794-801.
26. Mathew PJ, Fontes ML, Tudor CI, et al. James Ramsay et al. A Multicenter Risk Index for Atrial Fibrillation after Cardiac Surgery. *JAMA.* 2004;291:17201729.
27. Jakubová M1, Mitro P, Stančák B, Sabol F, Kolesár A, Cisarik P, Nagy V. The occurrence of postoperative atrial fibrillation according to different surgical settings in cardiac surgery patients. *Interact Cardiovasc Thorac Surg.* 2012 Dec;15(6):1007-12